

# COMMUNICATION ON COUNTING POLYNOMIALS FOR ALKANES

Lorentz JÄNTSCHI<sup>1</sup>, Carmen Elena STOENOIU<sup>1</sup>, Sorana Daniela BOLBOACĂ<sup>2</sup>

<sup>1</sup>Technical University of Cluj-Napoca, 15 Constantin Daicoviciu, 400020 Cluj-Napoca, Romania.

<http://lori.academicdirect.org/>, <http://carmen.academicdirect.ro/>

<sup>2</sup>„Iuliu Hațieganu“University of Medicine and Pharmacy, 13 Emil Isac, 400023 Cluj-Napoca, Romania <http://sorana.academicdirect.ro/>

## ABSTRACT

Henry's law constant of a sample of nonane isomers was modeled by using characteristic and counting polynomials. The characteristic polynomial and counting polynomials on the distance matrix, on the maximal fragments matrix, on the complement of maximal fragments matrix, and on the Szeged matrix were calculated for each compound and multi-varied models were identified and analyzed. Two multi-varied models, one with four and other with five variables, revealed to had estimation abilities. Both models used the characteristic and counting polynomials on Szeged matrix as criteria. The statistical characteristics of the models were analyzed and are presented. The obtained results shown that Henry's law constant of studied nonane isomers could be estimated by using characteristic polynomial and counting polynomial on Szeged matrix.

## INTRODUCTION

Graph theory, defines as the study of graphs that are mathematical structures used to model pair-wise relations between objects from a certain collection, was introduced by Leonhard Euler in 1736 [1]. Sub-graphs that result through applying of the matrix criteria of fragmentation are using in investigation of chemical structures and of relationships between structures and properties. Some criteria as characteristics polynomial, Szeged and Cluj matrices are used methods. Two new criteria that proved to had abilities in characterization of compounds properties were recently introduced and analyzed [2].

The aim of the present research was to investigate the abilities in estimation of Henry's law property of nonane isomers by using characteristic and counting polynomials.

## MATERIAL & METHOD

A sample of thirty-five compounds, representing the isomers of nonane (C9H20), where included into the study. The values of Henry's law constant (the ratio of the gaze phase concentration and the liquid phase concentration of a substance), expressed as [M/atm] unit, were taken from a previously reported research [3].

The Henry's law constant was modeled by using characteristic polynomial and counting polynomials. Five matrices were used in counting polynomials: characteristic polynomial (ChP), counting polynomial on the maximal fragments matrix (CMx), counting polynomial on the complementary of the maximal fragments matrix (CcM), and counting polynomial on the Szeged matrix (CSz). Comparisons between correlation coefficients obtained by different models were analyzed by using the Steiger's Z test [4] at a significance level of 5%.

## RESULTS

One out of thirty-five compounds (4-methyl-octane) was considered an outlier and was not included into the analysis. Two models proved to had abilities in estimation of the Henry's law constant for the nonane isomers.

- Model with four variables:

$$\hat{Y}_{KHL-4v} = 1100 - 1.23 \cdot 10^{-4} \cdot P_{CSz}(-2.83) + 0.1269 \cdot P_{ChP}(-1.72) + 2.73 \cdot 10^{-5} \cdot P_{ChP}(0.33) - 2.61 \cdot P_{ChP}(-9.36) \quad (1)$$

where  $\hat{Y}_{KHL-4v}$  is the estimated Henry's law constant by the model with four variable,  $P_{CSz}(X_i)$  is counting polynomial on the Szeged matrix and  $P_{ChP}(X_i)$  are characteristic polynomials.

- Model with five variables:

$$\hat{Y}_{KHL-5v} = 5669 - 3.36 \cdot 10^{-2} \cdot P_{CSz}(-3.07) + 1.80 \cdot 10^{-2} \cdot P_{CSz}(0.9) + 8.67 \cdot P_{ChP}(0.11) - 1.74 \cdot 10^{-2} \cdot P_{ChP}(1.83) - 3.70 \cdot 10^{-4} \cdot P_{ChP}(7.11) \quad (2)$$

where  $\hat{Y}_{KHL-5v}$  is the estimated Henry's law constant by the model with five variable,  $P_{CSz}(X_i)$  are counting polynomials on the Szeged matrix and  $P_{ChP}(X_i)$  are characteristic polynomials.

The statistical characteristics of the models are presented in table 1.

Table 1. Statistical characteristics of the models

Parameter	Model	
	Eq.(1)	Eq.(2)
Correlation coefficient (r)	0.9663	0.9705
95% confidence interval for (r)	[0.9329-0.9831]	[0.9411-0.9852]
Squared correlation coefficient ( $r^2$ )	0.9337	0.9418
Model significance	< 0.0001	< 0.0001

The graphical representation of the model obtained by Eq.(2) versus the measured Henry's law constant is shown in figure 1.

In order to test the null hypothesis that the correlation coefficient obtained by Eq.(1) is not different by the correlation coefficient obtained by Eq.(2), the Steiger's Z test was applied. The value of Z test was equal with 0.657 ( $p = 0.255$ ) showing us that there are not statistical difference between the correlation coefficients.




Figure 1. Estimated by Eq.(2) versus measured Henry's law constant for studied nonane isomers

## CONCLUSION

The characteristic polynomial and counting polynomial on Szeged matrix criteria revealed to be useful in modeling of Henry's law constant for studied nonane isomers. Both models (Eq.(1) and Eq.(2)) revealed to had abilities in estimation of the Henry's law constant. External and internal validations of the models are intended to be done in future research in order to characterize the validity and relevance of the obtained models.

## REFERENCES

- [1] Biggs, N.; Lloyd, E.; Wilson, R. Graph Theory. Oxford University Press (1986) 1736-1936.
- [2] Jäntschi, L.; Bolboacă, S.D. Counting Polynomials on Regular Structures, MATCH, Sent for publication.
- [3] Yaws, C. L.; Yang, H.-C. Henry's law constant for compound in water. In C.L. Yaws, editor, Thermodynamic and Physical Property Data, pp. 181–206. Gulf Publishing Company, Houston, TX, 1992.
- [4] Steiger, J.H.: Tests for comparing elements of a correlation matrix. Psychol. Bull. 87 (1980) 245-251.